

Version: 03 Revision date: 07/06/2023 Canvax Reagents, S.L.U.

Luis de Mercado Street, 19 Boecillo Technological Park 47151, Boecillo Valladolid, Spain.

Tlf: +34 983 54 85 63 info@canvaxbiotech.com

www.canvaxbiotech.com

1. Identification

Product name HigherPurity™ Soil DNA Isolation Kit

 Cat. No.
 AN0140 (50 rxn)

 Cat. No.
 AN0141 (100 rxn)

 Cat. No.
 AN0141-XL (250 rxn)

2. Description

HigherPurity™ Soil DNA Isolation Kit provides a simple and convenient technique to isolate high quality DNA from soil samples. Extraction is based in DNA ability to bind silica in the presence of high concentrations of chaotropic salts as guanidinium thiocyanate. Samples are rapidly and efficiently lysed by bead beating. The sample DNA is then bound to the surface of a silica membrane that is inside the spin column and washed and the bound DNA is then desorbed from the surface of the membrane. The inhibitors of the downstream PCR will be removed with the buffers system in this kit.

3. Kit Contents

Component	Quantity		
	AN0140 (50 rxn)	AN0141 (100 rxn)	AN0141-XL (250 rxn)
Lysis Solution 1 (LS1)	35 ml	65 ml	160 ml
Buffer A	12 ml	24 ml	60 ml
Inhibitor Removal Buffer (IR-Buffer)	10 ml	20 ml	40 ml
Buffer B	20 ml	40 ml	100 ml
Wash Buffer 1 * (WB1)	16 ml	32 ml	80 ml
EB Buffer	15 ml	30 ml	75 ml
CleanEasy™ MiniSpin Columns	50	100	250
Collection tube (2mL)	100	200	500
Bead Tube (with glass beads)	50	100	250
1.5 ml microcentrifuge tube	50	100	250

^{*}Add the volume ethanol (96%-100%) specified [Not included] to WB1 Buffer prior to initial use (see bottle label for volume). After ethanol has been added, mark the bottle to indicate that this step has been completed.

4. Storage specifications

All components can be stored at Room Temperature.



[!] Caution: Please wear gloves when using this product. Avoid all skin contact with kit reagents. In case of contact, wash thoroughly with water.



DATA SHEET

Version: 03 Revision date: 07/06/2023 Canvax Reagents, S.L.U.

Luis de Mercado Street, 19 Boecillo Technological Park 47151, Boecillo Valladolid, Spain.

Tlf: +34 983 54 85 63 info@canvaxbiotech.com

www.canvaxbiotech.com

5. Applications

All molecular biology applications, such as:

- Digestion with restriction enzymes.
- > Automated sequencing.
- > PCR template.
- Southern Blots.

6. Quality Certifications

Soil DNA Isolation Kit is tested for isolation of DNA from soil sample. The quantity and quality of purified DNA attend to:

- Ratio 260/280.
- Agarose gel electrophoresis.
- Digestion with restriction endonucleases

7. Further information

Product Use Limitations This product is developed, designed, and sold exclusively only for research purposes use. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or


animals.

Safety Information When working with chemicals, always wear a suitable lab coat, disposable gloves, and protective goggles. For more information, please consult the appropriate material safety data sheets (MSDS). These are available online in convenient and compact PDF format at www.canvaxbio.com where you can find, view, and print the MSDS for each CANVAX kit.

8. Assay Procedure

- 1. Transfer 0.25-1g (or 200 µl for liquid sample) of soil sample into Bead Tube and place on ice.
- 2. Add **0.6 mL of Lysis Solution 1 (LS1)**. Vortex for 5 minute at maximum speed. *Make sure that soil sample is homogenized completely.*
- 3. Incubate the sample at 70 °C for 10 minutes.

 The temperature can be increased to 95°C for isolation of DNA from gram positive bacteria.
- 4. Spin the tube to remove drops from the inside of the lid.
- 5. Cool down the sample on ice and add 200 μ l of Buffer A to the sample, mix well by vortexing. Incubate the sample on ice for 5 minutes.
- 6. Vortex for 10 seconds and centrifuge at full speed (14 000 rpm) for 5 minutes. Pipet the supernatant into a new 1.5 ml microcentrifuge tube (not provided) and discard the pellet. Measure the volume of the supernatant.
- 7. Add 1 volume of isopropanol (not provided) to the sample. Mix thoroughly by vortexing and centrifuge at full speed for 10 minute to pellet DNA.
- **8.** Carefully discard the supernatant and invert the tube on the paper towel for ~1 minute to drain any excess liquid from the pellet. *Do not disturb the pellet.*







DATA SHEET

Version: 03
Revision date: 07/06/2023

Canvax Reagents, S.L.U.

Luis de Mercado Street, 19 Boecillo Technological Park 47151, Boecillo Valladolid, Spain.

Tlf: +34 983 54 85 63 info@canvaxbiotech.com

www.canvaxbiotech.com

- 9. Add 200 µl of pre-heated Elution Buffer (65°C) and vortex to dissolve the DNA pellet completely.
- **10.** Add **100 µl of IR-Buffer** to the sample, mix well by vortexing. Incubate the sample at room temperature for 2 minutes. *IR-Buffer must be suspended completely by vigorously vortexing before every using.*
- **11.** Centrifuge at full speed for 3 minutes. Carefully pipet the supernatant into a new 1.5 ml microcentrifuge tube (not provided) and discard the pellet. *Measure the volume of the supernatant.*
- 12. Spin the tube to remove drops from the inside of the lid.
- **13.** Add 1 volume of **Buffer B** and 1 volume of ethanol (96%~100%). Mix thoroughly by vortexing.
- **14.** Assemble a spin column with one of the provided collection tubes. Apply the sample mixture onto the spin column. Close the cap and centrifuge at full speed for 1 minute. Discard the flow-through and and reuse the collection tube.
- **15.** Carefully open the spin column and add **750 µl Buffer WB1** (ethanol added). Close the cap and centrifuge at full speed for 1 min. Discard the flow-through and and reuse the collection tube.
- 16. Repeat step 15 for one more time.
- **17.** Centrifuge at full speed for an additional 3 min to dry the spin column. This step will avoid the residual liquid to inhibit subsequent enzymatic reactions.
- 18. Place the spin column into a new 1.5 mL microcentrifuge tube. Carefully open the spin column and Add 50-200 µl of water (60-70°C) to the membrane center. Water is non-provided. Close the cap and incubate for 1 min at room temperature, then centrifuge at full speed for 1 min to elute DNA.
- **19.** The purified genomic DNA can be stored at 2-8°C for a few days. For longer term storage, -20°C is recommended.

